

Regular Article

Nanoencapsulated deltamethrin combined with indoxacarb: An effective synergistic association against aphids

Marine Galloux,^{1,2} Guillaume Bastiat,² Corinne Lefrancois,¹ Véronique Apaire-Marchais^{1,*} and Caroline Deshayes¹


¹ Univ Angers, INRAE, SIFCIR, SFR QUASAV, F-49000 Angers, France

² Univ Angers, Inserm, CNRS, MINT, SFR ICAT, F-49000 Angers, France

(Received January 25, 2024; Accepted June 3, 2024)

Supplementary material

Widespread pesticide use for decades has caused environmental damage, biodiversity loss, serious human and animal health problems, and resistance to insecticides. Innovative strategies are needed to reduce treatment doses in pest management and to overcome insecticide resistance. In the present study, combinations of indoxacarb, an oxadiazine insecticide, with sublethal concentrations of deltamethrin encapsulated in lipid nanocapsules, have been tested on the crop pest *Acyrtosiphon pisum*. *In vivo* toxicological tests on *A. pisum* larvae have shown a synergistic effect of nanoencapsulated deltamethrin with a low dose of indoxacarb. Furthermore, the stability of deltamethrin nanoparticles has been demonstrated *in vitro* under different mimicking environmental conditions. In parallel, the integrity and stability of lipid nanoparticles in the digestive system of aphid larvae over time have been observed by Förster Resonance Energy Transfer (FRET) imaging. Thus, the deltamethrin nanocapsules/indoxacarb synergistic association is promising for the development of future formulations against pest insects to reduce insecticide doses.



Keywords: indoxacarb, *Acyrtosiphon pisum*, nanoencapsulated deltamethrin, synergistic association, pest control.

Introduction

Pests destroy up to 40% of global crop production and cause US \$220 billion in trade losses each year according to the United Nations and invasive insects cost a minimum of US \$70 billion per year globally.^{1,2} Among pest insects, the aphids especially pea aphid *Acyrtosiphon pisum*, a key pest of leguminous and pulse crops worldwide³ can cause serious economic damage and yield losses. This sap-sucking insect has a wide host range including fava bean, lupin, alfalfa, lentil, chickpea, grass pea, and

field pea. *A. pisum* as phloem-feeders can injure crops directly by consuming the sap of plants and indirectly by vectoring over 30 phytopathogenic viruses, including cucumber mosaic virus, pea enation mosaic virus, and bean leafroll virus.^{4,5} Aphids have a complex life cycle, including a succession of parthenogenetic generations during spring and summer and a single sexual generation in autumn.^{6,7} This mode of reproduction together with i) very prolific parthenogenetic adult females capable of producing 50 to 100 nymphs, ii) short generation time of aphids as well as iii) their remarkable ability adaptation to environmental conditions contribute to the rapid growth of aphid populations and make the control of these crop pests difficult.^{3,7,8} Currently, among strategies developed to fight against aphids, chemical insecticides are still primarily used. However, the development of resistance to insecticides must be considered in management strategy for aphid control.^{3,4,7,9}

Since decades, Integrative Pest Management (IPM) and Insecticide Resistance Management (IRM) programs have been developed to evaluate the proper insect pests control methods

* To whom correspondence should be addressed.

E-mail: veronique.marchais@univ-angers.fr

Published online July 9, 2024

© Pesticide Science Society of Japan 2024. This is an open access article distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

both for effectiveness and risk. According to the Food and Agriculture Organization of the United Nations, IPM is defined as “an ecosystem approach to crop production and protection that combines different management strategies and practices to grow healthy crops and minimize the use of pesticides”.¹⁰ Many networks like PAN Europe or ENDURE have been created to contribute to IPM strategies implementation.^{11,12} However, a gap exists between IPM concepts and practices in the fields.¹³ Indeed, despite i) the increasing development of insecticide resistances, ii) the rising number of restrictions regarding the use of agrochemicals with high health risks, iii) the implementation of laws or national programs to reduce the adverse effects on the environment, the beneficial insects and on the human health, current pest control strategies still rely on the use of insecticides.^{14–16} These problems reinforce the need to develop more efficient strategies allowing a decrease in the amount of active ingredients in the formulations used for insecticide treatments.

Among innovative strategies to reduce doses of insecticides treatment, it has previously been shown that sublethal concentration of deltamethrin was able to potentiate the effect of indoxacarb, an oxadiazine insecticide, on the cockroach *Periplaneta americana*.¹⁷ Indoxacarb is a pro-insecticide that undergoes bioactivation to the more effective *N*-decarbomethoxyllated indoxacarb (DCJW) by esterases and amidases in insects.^{18,19} This insecticide is part of the group 22A in the mode of action classification table of insecticide resistance action committee IRAC²⁰ as the DCJW metabolite is a highly potent voltage-gated sodium channel blocker.^{21,22} In this strategy, deltamethrin produces an increase of intracellular calcium concentration in isolated insect neurons through the reverse Na/Ca exchanger leading to activation of calcium-dependent intracellular signalling pathways. This activation induces a modification in the voltage-gated sodium channel conformational states, leading to increase its sensitization to DCJW, thus optimizing the efficacy of the treatment. Furthermore, to protect deltamethrin from insect detoxification enzymes like esterases and cytochrome P450 monooxygenases, this molecule was nanoencapsulated in lipid nanocapsules. This was an essential first step to optimize the synergistic association when combined with indoxacarb.¹⁷ Because this previous study demonstrated that nanoencapsulated deltamethrin combined with indoxacarb increased cockroaches mortality rate at lower concentrations compared to indoxacarb used alone, we decided to adapt and to test such strategy on aphid, *A. pisum*. The pro-insecticide indoxacarb has been designated by the EPA to be a “reduced-risk” pesticide and is used against a broad spectrum of insect pests. This insecticide is therefore compatible with IPM programs^{23–25} and it has been reported to be effective on aphids under laboratory conditions.^{26,27} Here, toxicological tests on *A. pisum* larvae have demonstrated the efficacy of the deltamethrin nanocapsules/indoxacarb association. In parallel, Förster Resonance Energy Transfer (FRET) imaging has been used for monitoring *in vivo* the integrity of lipid nanoparticles, showing a stability and storage in aphid larvae digestive system over time. Furthermore, *in vitro* experiments have demonstrated a high

stability of deltamethrin-nanoparticles against various environmental-mimicking conditions, making this association a promising strategy in pest management.

Materials and methods

1. Chemicals

Deltamethrin (Sigma-Aldrich, Saint-Quentin-Fallavier, France) was diluted in dimethylsulfoxide (DMSO) (Sigma-Aldrich) to obtain a stock solution at 10^{-2} M and stored at -20°C . The insecticide indoxacarb (Avaunt® 15% Emulsion Concentrate (EC)) was supplied by DuPont™ (Wilmington, Delaware, USA). Piperonyl butoxide (PBO) (Sigma-Aldrich) was prepared in DMSO to obtain a stock solution at 500 mg/mL stored at 4°C . Lipophilic carbocyanine dyes 1,1'-dioctadecyl-3,3',3'-tetramethylindodicarbocyanine 4-chlorobenzenesulfonate (DiD) and 1,1'-dioctadecyl-3,3',3'-tetramethylindocarbocyanine perchlorate (DiI) were supplied by Molecular probes® (Eugene, OR, U.S.A.) and stored at 4°C .

2. Insects

Experiments were performed on the pea aphids, *Acyrtosiphon pisum* strain LSR1.²⁸ They were generously provided by INRAE-Rennes IGEPP. Unwinged parthenogenetic females were reared on faba bean (*Vicia fabae*) plants under laboratory conditions at 22°C and a photoperiod of 16:8 hr (L:D). Under these conditions, aphids reproduced *via* viviparous parthenogenesis as clonal females, and newborn larvae became adults after four molts.

3. In vivo toxicological studies

All the experiments were conducted with an artificial diet bioassay as previously developed.^{29,30} According to the method, artificial diet was prepared and stored at -20°C for a period up to 6 months. The artificial diet was used to prepare all compound dilutions. Tests were performed using 100–375 larvae (for $n=4$ to 15) per concentration of insecticide. In brief, adults were put on a feeding apparatus containing 200 μL of artificial diet at day 0. At day 1, 25 first-instar nymphs were placed to 200 μL of freshly prepared diet containing molecules alone or in combination. Concentrations ranging from 10^{-3} to 10^{-8} M for indoxacarb and from 10^{-4} to 10^{-8} M for deltamethrin were tested. For PBO, a sublethal concentration of 1 mg/mL was used. The mortality was observed after 48 hr (day 3) and 72 hr (day 4). Aphids that were unable to walk were considered dead and were removed.

Corrected mortality percentages were calculated using Henderson Tilton's formula after 48 hr and 72 hr of insecticide exposure.³¹ The Model Deviation Ratio (MDR) (observed toxicity/predicted toxicity) was calculated to define the nature of the association.³² MDR values <0.7 mean “antagonistic effect”, results between 0.7 and 1.3 mean “absence of effect” and results >1.3 mean “synergistic effect”.

4. Formulation and characterization of lipid nanocapsules

4.1. Preparation of lipid nanocapsules, deltamethrin-loaded lipid nanocapsules and dye-loaded lipid nanocapsules

Lipid nanocapsules (LNCs) were performed following a previously developed phase inversion-based method.³³⁾ Briefly, 75 mg of Lipoid® 75-3 (Lipoid GmbH, Ludwigshafen, Germany) and 846 mg of Kolliphor® HS 15 (BASF, Ludwigshafen, Germany) (surfactants), 1028 mg of Labrafac® WL 1349 (oil phase) (Gattefossé S.A., Saint-Priest, France), 89 mg of NaCl (Prolabo, Fontenay-Sous-Bois, France) and 2962 mg of de-ionized water (Milli® Q system, Millipore, Billerica, MA, USA) (aqueous phase) were precisely weighed.³⁴⁾ Under continuous agitation (magnetic stirring), the mix was heated up to 95°C and cooled down to 50°C three times. During the last cooling, 5 mL of de-ionized cold water (at 4°C) were quickly added at about 75°C corresponding to the transition between the oil-in-water (low temperature) and water-in-oil (high temperature) emulsions, to stop the emulsion process, leading to the suspension of LNCs. For deltamethrin-loaded LNCs (deltamethrin-LNCs) formulation, deltamethrin was first dissolved in Labrafac® WL 1349 at a concentration of 1 mg/g Labrafac® WL 1349 (w/w), before adding other constituents.¹⁷⁾ Before dye-loaded LNCs formulation, 4-chlorobenzenesulfonate and perchlorate ions for DiD and DiI, respectively, were replaced by tetraphenylborate (TPB) ion to improve dye solubility in Labrafac® WL 1349 and consequently the dye loading capacity.³⁵⁾ Then, these modified dyes were solubilized in Labrafac® WL 1349 before formulation process to achieve DiI/TPB-loaded LNCs (DiI/TPB at 1% (w/w) in Labrafac® WL 1349) (DiI/TPB-LNCs), DiD/TPB-loaded LNCs (DiD/TPB at 1% (w/w) in Labrafac® WL 1349) (DiD/TPB-LNCs), and both DiI/TPB and DiD/TPB-loaded LNCs (both DiI/TPB and DiD/TPB at 1% (w/w) in Labrafac® WL 1349) (FRET-LNCs). LNCs, deltamethrin-LNCs, DiI/TPB-LNCs, DiD/TPB-LNCs and FRET-loaded LNCs suspensions were stored at 4°C.

4.2. Characterization of lipid nanocapsules and deltamethrin-loaded lipid nanocapsules

LNCs and deltamethrin-LNCs were diluted by factor 60 (v/v) in de-ionized water and size distribution, *i.e.*, hydrodynamic diameter (Z-ave), polydispersity index (PDI) and Zeta potential (ZP) were measured using a Zetasizer® Nano ZS (Malvern Panalytical Ltd., Worcestershire, UK) at 25°C. The instrument is equipped by a Helium–Neon laser, 4 mW, with an output wavelength of 633 nm and a scatter angle fixed at 173°. The correlation functions were fitted using an exponential fit (Cumulant approach) for Z-ave and PDI determinations. Smoluchowski approach was used to evaluate the electrophoretic mobility to calculate ZP. The encapsulation efficiency of deltamethrin in LNCs was evaluated by chromatography, as previously described.¹⁷⁾

4.3. LNC stability

The stability of deltamethrin-LNCs was followed over time up to 14 days depending on different settings: temperature (4, 20 and 37°C), pH (4, 6, 8, 10) and brightness (photoperiod of 16:8 hr (L:D)) or darkness, UV 365 nm (24 hr). Deltamethrin-LNCs were diluted to 7/10 in de-ionized water and four parameters

(Z-ave, PDI, ZP and encapsulation efficiency) were characterized as described above.

5. Statistical analysis

Data were analyzed with GraphPad Prism version 5.00 (GraphPad software, Inc.) and displayed as mean and S.E.M. (Standard error of the mean). Values are means of at least three independent experiments. Between-group comparisons were assessed by nonparametric Mann–Whitney *U*-test or Kruskal–Wallis test followed by *post-hoc* Dunn test for pairwise comparison. A $p < 0.05$ was considered statistically significant.

6. In vivo imaging

6.1. Monitoring of LNCs by FRET

FRET methodology was used to monitoring the integrity of LNCs *in vivo*.³⁶⁾ Studies were realized on first instar larvae (aged 0–24 hr). Dye-loaded LNCs were diluted with artificial diet to obtain a concentration of 850 and 85 µg/mL. Ten larvae were placed in diet cages and fed with different tests' conditions (DiI-LNCs, DiD-LNCs and FRET-LNCs) for 5 or 24 hr. Then tests' conditions were removed and replaced by artificial diet for 24 hr to 7 days. At the end of each experiment, larvae were frozen at –20°C.

6.2. Fluorescence analyzing

Fluorescence analyzing was performed at Plateau Technique Mutualisé d'IMagerie Cellulaire (PTM IMAC) (SFR 4207 Quasav, Angers, France). Larvae having previously ingested dye-loaded LNCs were placed on slide (Hecht Karl™, Sondheim, Germany), and recovered by a coverslip (Thermo Fisher Scientific Inc., Illkirch, France). Larvae were thus placed under fluorescence microscope (NIKON, Ti-E A1, France), at a CFI Plan Fluor 10X objective (NA 0.30, WD 16.0 mm). Adequate excitation and emission wavelength couples ($\lambda_{ex}/\lambda_{em}$) were used to analyze fluorescence. These couples were 561/570–620 for DiI, 640/663–738 nm for DiD and 561/663–738 for FRET. Data obtained were compared each other with application of specific settings common to all samples (excitation time 10 msec, laser powerful 2%) with the software NIS-Element Arv 5.21.03 (NIKON Corporation, Tokyo, Japan).

Results and discussion

1. Studies of the toxicity of indoxacarb and deltamethrin used alone and in combination on *A. pisum* larvae

To evaluate the potential synergy between deltamethrin and indoxacarb *in vivo* on *A. pisum* larvae, we first investigated the effect of each molecule alone on the mortality rate. The toxicological effects of deltamethrin or indoxacarb upon these sucking insects were studied by using an artificial diet system as previously described.²⁹⁾ The average corrected percentages of mortality were plotted against the noncumulative concentrations of indoxacarb or deltamethrin after 48 hr and 72 hr ingestion. The mortality rate after 24 hr ingestion was not investigated as preliminary experiments showed absence of larval mortality after ingestion of indoxacarb. The sigmoid curves corresponded to

Table 1. Parameters of the concentration–response curves established for indoxacarb and deltamethrin in *A. pisum* larvae after 48 hr and 72 hr ingestion

	Indoxacarb		Deltamethrin	
	48 hr	72 hr	48 hr	72 hr
LC ₅₀ (M)	1.1 10 ⁻⁵ M	6.2 10 ⁻⁶ M	7.8 10 ⁻⁶ M	6.3 10 ⁻⁶ M
Hill slope	1.93	5.82	1.50	2.67
R ²	0.95	0.96	0.91	0.88

the fit according to the Hill equation (Table 1). The concentration–response curve established with indoxacarb after 72 hr post-ingestion was steeper than the curve at 48 hr after ingestion. The corresponding slope parameters were 1.93 and 5.82 at 48 hr and 72 hr, respectively (Fig. 1A, Table 1). This variation may reflect the bioactivation of indoxacarb in its active metabolite DCJW. The increase of the Hill slope parameter at 72 hr could therefore be due to a higher quantity of DCJW, which has a better binding affinity to the voltage-gated sodium channel compared to indoxacarb. The maximum corrected percentage of mortality was 100% for 48 hr ($n=3$) and 72 hr ($n=4$) after 10⁻³ M indoxacarb ingestion, respectively. The threshold concentration inducing very low mortality (<10%) was 3×10⁻⁶ M at 48 hr and 72 hr (Fig. 1A). The LC₅₀ values estimated for indoxacarb were 1.1×10⁻⁵ M and 6.2×10⁻⁶ M at 48 hr and 72 hr after ingestion, respectively (Table 1).

For deltamethrin, the Hill slope was slightly increased for the sigmoid curve obtained after 72 hr ingestion (Hill slope 2.67) compared to the concentration–response curve at 48 hr (Hill slope 1.50) (Fig. 1B, Table 1). The maximum corrected mortality rates were 93±3% ($n=6$) and 99±1% ($n=6$) at 48 hr and 72 hr after ingestion, respectively. The LC₅₀ values estimated for deltamethrin were 7.8×10⁻⁶ M and 6.3×10⁻⁶ M at 48 hr and 72 hr after ingestion, respectively. The 48 hr post-ingestion LC₅₀ of deltamethrin in *A. pisum* had been previously evaluated to

3.1×10⁻⁵ M, which is in the same concentration range compared to our results.³⁷⁾

This set of experiments allowed to determine the lower concentrations of indoxacarb and deltamethrin to be tested for the synergy study between both compounds. Indeed, in our strategy, deltamethrin was used at concentrations that induce the lowest lethal effect, it must therefore be used at concentrations less than or equal to 10⁻⁷ M.

Thus aphid larvae were fed with various concentrations of indoxacarb (10⁻⁷ to 10⁻⁵ M) alone and associated with deltamethrin (10⁻⁷ M). The percentages of corrected mortality after 48 hr and 72 hr ingestion are illustrated in Fig. 2. No significant difference was observed between mortality rates with indoxacarb alone or associated with deltamethrin 10⁻⁷ M at 48 hr or 72 hr after ingestion.

2. Physico-chemical in vitro stability of deltamethrin-LNCs

The absence of synergy between the deltamethrin and indoxacarb probably reflects the rapid metabolization of deltamethrin by detoxification enzymes of aphid larvae.^{38,39)} It has been previously shown that encapsulation of deltamethrin in lipid nanocapsules (LNCs) protects deltamethrin from enzymatic degradation allowing the optimization of the synergistic association.¹⁷⁾ Since few years, nanotechnology research is raising interest in the agricultural sector as nano-based pesticide formulations have several advantages such as increased efficacy, controlled release or reduced environmental impacts.^{40–45)} Indeed various classes of insecticides have been already nanoformulated like pyrethroids (deltamethrin, β -cypermethrin), benzoylphenyl urea (novaluron) or avermectines (abamectine).^{41,44)} Among these nano-insecticides, various types of nanosystems have been developed such as nanoemulsion, nanospheres, nanomicelles, nanosuspensions or nanocapsules. In the latter the insecticides are encapsulated in a core-shell structural vesicular system usually composed of biodegradable polymers (*i.e.*, chitosan, poly-




Fig. 1. Toxicity of indoxacarb and deltamethrin on *A. pisum* larvae. Aphids were fed orally with artificial diet containing indoxacarb (A) ranging from 10⁻⁸ to 10⁻³ M or deltamethrin (B) ranging from 10⁻⁸ to 10⁻⁴ M. Mortality rate was assessed 48 hr and 72 hr after intoxication. Smooth lines represent the fit through the mean data according to the Hill equation. Data are means±S.E.M. ($n=4-15$).




Fig. 2. Toxicity of deltamethrin/indoxacarb association on *A. pisum* larvae. Comparative mortality rates produced by indoxacarb (10^{-7} to 10^{-5} M) tested alone and indoxacarb associated with deltamethrin at 10^{-7} M at 48 hr (A) and 72 hr (B). Data are means \pm S.E.M. ($n=4-13$). ns: not significant $p>0.05$.

ethylene glycol, poly(lactic-co-glycolic acid)) whom slow degradation in the environment increases the chemical stability.^{17,45,46} Consequently, we first studied the physico-chemical stability of deltamethrin-loaded nanocapsules (deltamethrin-LNCs) under various conditions mimicking environmental factors affecting release. The size distributions of deltamethrin-LNCs were measured *in vitro* after incubation at different conditions of temperature (4, 20 and 37°C), pH (4, 6, 8, 10) and brightness or darkness, (photoperiod L:D), UV 365 nm (24 hr). All conditions tested were compared to reference values obtained at 20°C, pH 6, in the presence of light at time 0. Regardless of all the parameters tested, encapsulation values remain constant over time, from 24 hr to 14 days, i.e., Z-ave=56–64 nm, PdI=0.05–0.12, ZP=−13–−6 mV and deltamethrin encapsulation rate=92–113% (Table 2). The deltamethrin-LNCs did not change at acidic or basic pH values and were not degraded in different conditions, suggesting that the deltamethrin still remained encapsulated within the LNCs and protected in the external environment before ingestion by the insects.

3. Lipid nanocapsules fate in *A. pisum* larvae

We then focused on the *in vivo* stability and fate of the nanocapsules. To carry out this study, control experiments have been performed to estimate the toxicity of empty LNCs on *A. pisum* larvae. Aphids were fed with artificial diet containing different concentrations of LNCs (from 8.5 to 8500 μ g/mL). The highest tested concentration corresponded to the maximal concentration that could be obtained in the experimental system. At the concentration of 8500 μ g/mL, LNCs were toxic for aphid larvae with corrected mortality of $64 \pm 4\%$ at 48 hr and $91 \pm 7\%$ at 72 hr (Fig. 3). However, no effect on mortality rates was observed for lower LNCs concentrations and *in vivo* studies were then performed with LNCs at the maximal concentration of 850 μ g/mL.

The fate and integrity of LNCs in *A. pisum* larvae were monitored by FRET (Förster resonance energy transfer) imaging. Co-encapsulation of two fluorescent probes (DiI, FRET donor, and DiD, FRET acceptor) in LNCs (FRET-LNCs) enables to monitor the nanocapsules stability *in vivo*.³⁵

Before tracking the LNCs integrity in *A. pisum* larvae over

time, we had to define experimental conditions for which a FRET signal was obtained after FRET-LNCs ingestion. Different ingestion times (5 or 24 hr) and concentrations (850 or 85 μ g/mL) of FRET-LNCs were tested. Analysis by fluorescent microscopy in whole insects has shown a stronger FRET signal after 24 hr ingestion of FRET-LNCs at 850 μ g/mL compared to shorter time or lower concentration of FRET-LNCs (Supplementary Fig. 1). LNCs integrity was thus tracked *in vivo* over 8 days after 24 hr ingestion of FRET-LNCs at 850 μ g/mL. As shown on Fig. 4, a FRET signal has been observed over 8 days with a maximum intensity around 72–96 hr post-ingestion. This FRET signal was correlated with a weak signal of DiI, the donor molecule, indicating a decrease of the donor fluorescence intensity due to energy transfer to acceptor molecule. This signal was localized in a part of the abdomen of aphid larvae. The gut anatomy of *A. pisum* is very simple with a stomach followed by an intestine without diverticula.^{47,48} Based on our fluorescent microscopy observations, the FRET-LNCs seemed to be stocked in the stomach. At 72 hr post-ingestion, a FRET signal was observed in the whole abdomen, reflecting a probable diffusion of FRET-LNCs in *A. pisum* larvae. The presence of FRET signal indicated an integrity of LNCs over time. One may hypothesized that LNCs were dissociated in stomach, thus releasing the fluorescent probes, but that the latter remained sufficiently close for the transfer of energy. Two control experiments have been performed, aphid larvae have ingested a mix of free fluorescent probes (DiI and DiD) or a mix of DiI-LNCs and DiD-LNCs. No fluorescent signal was observed after 24 hr post-ingestion of free probes, indicating that even these probes were released, they were not stocked in stomach and capable of energy transfer (Supplementary Fig. 2). Control experiment with ingestion of DiI-LNCs/DiD-LNCs provided confirmation that full LNCs are concentrated in stomach after 24 hr post-ingestion, as FRET signal was observed, meaning that these LNCs were close enough to have energy transfer from DiI-LNCs to DiD-LNCs (Supplementary Fig. 2). Our results indicated that LNCs were stable over time in *A. pisum*. The integrity of LNCs *in vivo* is an important point for our strategy, as the deltamethrin needs to remain protected from enzyme degradation to allow time for the

Table 2. *In vitro* stability of deltamethrin-LNCs over time as a function of pH, temperature and light

a. Particle size (Z-ave) (nm)								
Experimental conditions			Time (days)					
Temp.	pH	Light	0	1	2	3	7	14
20°C	6	+	56±2	59±3	59±3	58±1	59±1	59±1
	4			57±2	59±4	58±1	59±1	59±1
	8			57±2	58±1	59±3	58±2	60±1
	10			57±1	57±1	59±2	59±2	62±5
4°C				57±1	57±1	57±1	58±1	59±1
37°C				57±1	57±1	57±1	58±1	64±11
		—		56±2	60±5	58±1	59±6	59±3
		UV		57±1				
b. Polydispersity index (PdI)								
Experimental conditions			Time (days)					
Temp.	pH	Light	0	1	2	3	7	14
20°C	6	+	0.06±0.02	0.10±0.07	0.10±0.08	0.11±0.07	0.07±0.03	0.07±0.01
	4			0.06±0.01	0.10±0.11	0.06±0.01	0.06±0.02	0.07±0.01
	8			0.07±0.04	0.08±0.03	0.10±0.09	0.06±0.01	0.08±0.02
	10			0.06±0.01	0.06±0.01	0.11±0.09	0.09±0.05	0.11±0.08
4°C				0.09±0.04	0.06±0.02	0.07±0.01	0.07±0.01	0.07±0.01
37°C				0.08±0.04	0.06±0.01	0.08±0.03	0.06±0.02	0.12±0.13
		—		0.06±0.01	0.10±0.08	0.08±0.04	0.09±0.08	0.09±0.07
		UV		0.05±0.02				
c. Zeta potential (ZP) (mV)								
Experimental conditions			Time (days)					
Temp.	pH	Light	0	1	2	3	7	14
20°C	6	+	−12±5	−9±2	−8±1	−8±2	−9±3	−13±4
	4			−9±2	−9±1	−8±1	−9±2	−10±3
	8			−8±1	−7±1	−8±2	−8±2	−8±2
	10			−10±2	−9±2	−8±2	−10±4	−8±3
4°C				−8±1	−7±1	−7±2	−7±2	−6±2
37°C				−7±0	−9±2	−8±2	−7±2	−10±4
		—		−9±2	−8±1	−9±1	−8±2	−10±5
		UV		−9±1				
d. Deltamethrin encapsulation rate (%)								
Experimental conditions			Time (days)					
Temp.	pH	Light	0	1	2	3	7	14
20°C	6	+	105±3	106±3	104±7	109±4	107±5	104±6
	4			103±2	101±5	102±4	104±7	100±3
	8			101±4	102±5	101±6	94±6	92±9
	10			105±1	101±4	105±1	104±3	98±10
4°C				107±2	106±5	104±8	105±4	107±2
37°C				108±4	107±4	107±3	107±4	113±5
		—		110±2	109±4	108±3	109±4	112±4
		UV		105±2				

Four parameters have been studied: (a) particle size (Z-ave), (b) polydispersity index (PdI), (c) Zeta potential (ZP) and (d) deltamethrin encapsulation rate. $n=4$, mean±S.D.




Fig. 3. Toxicity of LNCs on *A. pisum* larvae. Comparative mortality rates produced by different concentrations (8.5 to 8500 µg/mL) of LNCs at 48 hr (A) and 72 hr (B). Data are means \pm S.E.M. ($n=4-11$).




Fig. 4. Lipid nanocapsules fate in *A. pisum* larvae. Fluorescence microscopy analysis over time in *A. pisum* larvae after 24 hr ingestion of FRET-LNCs at 850 µg/mL. FRET and DiI signals were observed using excitation wavelength of 561 nm. The excitation wavelength for acceptor DiD was 640 nm. For FRET and DiD signals, the emission wavelengths were between 663 and 738 nm. The emission wavelengths of DiI signal were between 570 and 620 nm. All sets of images were acquired with the same conditions. Scale bar: 500 µm.

indoxacarb to be bio-activated in DCJW.

4. Toxicity of deltamethrin-loaded nanocapsules associated with indoxacarb on *A. pisum* larvae

Results on deltamethrin-loaded LNCs stability under various

physico-chemical conditions and the fact that the LNCs remain intact *in vivo* over time in *A. pisum* indicated that deltamethrin-LNCs could be an added value in the crop protection.

Thereafter, the impact of the encapsulation of deltamethrin on *A. pisum* toxicity was investigated. The mortality rates of




Fig. 5. Impact of encapsulation on deltamethrin (deltamethrin-LNCs) toxicity on *A. pisum* larvae. Comparative mortality rates produced by deltamethrin alone and deltamethrin-LNCs at 10^{-8} M and 10^{-7} M at 48 hr (A) and 72 hr (B). Data are means \pm S.E.M. ($n=4-14$). ns: not significant $p > 0.05$; *** $p < 0.001$.

two concentrations of encapsulated deltamethrin (10^{-8} M and 10^{-7} M) were compared to mortality rates obtained with free deltamethrin (Fig. 5). For both concentrations of deltamethrin (10^{-7} M and 10^{-8} M), no significant difference was observed between encapsulated deltamethrin and non-encapsulated deltamethrin.

To study the efficacy of the association deltamethrin-LNCs/indoxacarb, aphids were fed with deltamethrin-LNCs at 10^{-8} M or 10^{-7} M of deltamethrin associated with various concentrations of indoxacarb (from 10^{-7} to 10^{-5} M). As no synergistic effect has been observed with associations of indoxacarb with deltamethrin-LNCs 10^{-8} M (data not shown), only comparative mortality rates with deltamethrin-LNCs at 10^{-7} M estimated at 48 hr and 72 hr are illustrated in Fig. 6. At 48 hr, significant differences were observed between corrected mortality of indoxacarb alone and deltamethrin-LNCs 10^{-7} M/indoxacarb for three concentrations of indoxacarb. Indeed, the mortality rate increased from 3 ± 2 to $34 \pm 5\%$ at 3×10^{-6} M ($U(6,6)=0.0$, $p=0.0048$), from 20 ± 5 to $50 \pm 12\%$ at 5×10^{-6} M ($U(6,4)=2.0$, $p=0.0381$) and from 49 ± 3 to $87 \pm 6\%$ at 10^{-5} M ($U(7,12)=6.0$, $p=0.0027$) (Fig. 6A). At 72 hr, significant differences in mortality rates were observed for concentrations of indoxacarb from 10^{-6} to 10^{-5} M. At this time, the corrected mortality increased

from 6 ± 1 to $37 \pm 7\%$ at 10^{-6} M ($U(13,16)=0.0$, $p=0.004$), from 11 ± 5 to $94 \pm 3\%$ at 3×10^{-6} M ($U(5,6)=0.0$, $p=0.0080$), from 28 ± 5 to $93 \pm 4\%$ at 5×10^{-6} M ($U(5,4)=0.0$, $p=0.0159$) and from 92 ± 3 to $99 \pm 1\%$ at 10^{-5} M ($U(8,12)=27.0$, $p=0.0377$) (Fig. 6B). The model deviation ratio (MDR) was calculated to define the nature of these associations (Table 3).^{32,49,50} At 48 hr, the MDR values were higher than 1.3 for three deltamethrin-LNCs/indoxacarb combinations meaning a synergistic effect. Indeed, the MDR values were 3.0, 1.8 and 1.4 for the deltamethrin-LNCs 10^{-7} M associated with indoxacarb at 3×10^{-6} M, 5×10^{-6} M and 10^{-5} M, respectively. At 72 hr, synergistic effect was observed with deltamethrin-LNCs 10^{-7} M associated with indoxacarb at 10^{-6} M, 3×10^{-6} M and 5×10^{-6} M (MDR values of 2.2, 2.9 and 2.8, respectively). The best synergistic effect was obtained with deltamethrin-LNCs at 10^{-7} M combined with indoxacarb at 3×10^{-6} M.

This synergy was not observed when deltamethrin was not encapsulated (Fig. 2). To ensure that potentialization of indoxacarb activity was due to encapsulated deltamethrin, control experiment was performed with the association indoxacarb 3×10^{-6} M and LNCs alone ($85 \mu\text{g/mL}$). The results indicated a higher mortality at 48 hr ($20 \pm 4\%$) ($Z(6.9)=2.574$, $p=0.0604$) and at 72 hr ($44 \pm 2\%$) ($Z(5.9)=2.296$, $p=0.1300$) compared to




Fig. 6. Toxicity of deltamethrin-LNCs/indoxacarb association on *A. pisum* larvae. Comparative mortality rates produced by indoxacarb (10^{-7} to 10^{-5} M) tested alone and indoxacarb associated with deltamethrin-LNCs 10^{-7} M at 48 hr (A) and 72 hr (B). Data are means \pm S.E.M. ($n=4-13$). ns: not significant $p > 0.05$; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Table 3. Model Deviation Ratio (MDR) values used to determine the nature of deltamethrin-LNCs/indoxacarb associations at 48 hr and 72 hr in *A. pisum*. MDR<0.7=antagonist effect; 0.7<MDR<1.3=absence of effect; MDR>1.3=synergistic effect. na, not applicable

	MDR	
	48 hr	72 hr
Deltamethrin-LNCs 10^{-7} M/indoxacarb 10^{-7} M	na	na
Deltamethrin-LNCs 10^{-7} M/indoxacarb 10^{-6} M	na	2.2
Deltamethrin-LNCs 10^{-7} M/indoxacarb 3×10^{-6} M	3.0	2.9
Deltamethrin-LNCs 10^{-7} M/indoxacarb 5×10^{-6} M	1.8	2.8
Deltamethrin-LNCs 10^{-7} M/indoxacarb 10^{-5} M	1.4	0.9

indoxacarb alone ($3 \pm 2\%$ and $11 \pm 5\%$ at 48 hr and 72 hr, respectively). One may hypothesize that this slight synergistic effect is due to LNCs which act as synergistic agent or to an interaction between indoxacarb and LNC which potentiates indoxacarb activity in insect cells (*i.e.*, increased local concentration of the insecticide). However, the potentialization of indoxacarb by LNCs alone did not reach the synergistic effect obtained with deltamethrin-LNCs at 10^{-7} M combined with indoxacarb at 3×10^{-6} M ($94 \pm 3\%$) indicating that the synergy is due to deltamethrin encapsulated in LNCs. Indeed, indoxacarb needs to be metabolized to DCJW by esterase/amidase enzymes. This metabolite is a potent voltage-gated sodium channel blocker.²²⁾ As previously shown in *Periplaneta americana*,¹⁷⁾ the deltamethrin may modulate voltage-gated sodium channels in aphids and increase intracellular sodium concentration, which could activate the Na/Ca exchanger in reverse mode. The voltage-gated sodium channels may thus be more sensitive to lower concentration of DCJW due to the elevation of intracellular Ca^{2+} concentration. However, an esterase-induced enzymatic degradation could occur on deltamethrin.¹⁷⁾ So nanoencapsulation of deltamethrin could protect its *in vivo* degradation allowing sufficient time to bioactivate indoxacarb in DCJW. This synergistic effect may be due to deltamethrin protection but also to its sustained release. Indeed, a synergistic effect was observed since 48 hr post ingestion while FRET experiments showed LNCs integrity and storage in stomach of *A. pisum* larvae until 8 days (Fig. 4). This combination of effects: protection against degradation enzymes, effective delivery or sustained release would allow a synergistic effect with the indoxacarb and reduction of insecticide concentrations, which corresponds to national guidelines.¹⁰⁾ Encapsulated deltamethrin could be associated with different other insecticides acting on voltage-gated sodium channel (*i.e.*, pyrethroids). The encapsulation strategy could improve the efficacy of synergistic agent associated with various insecticides.^{51–53)}

5. Comparative studies of the toxicity between deltamethrin-LNCs/indoxacarb and deltamethrin/indoxacarb associated with PBO

The toxicity of the deltamethrin-LNCs 10^{-7} M/indoxacarb 3×10^{-6} M association was then compared to the toxicity of deltamethrin 10^{-7} M/indoxacarb 3×10^{-6} M associated with the cur-

rent used synergist piperonyl butoxide (PBO). This experiment enabled to evaluate the *in vivo* protection of deltamethrin from enzymes degradation by nanocapsules *versus* PBO. Indeed PBO is known to inhibit the activity of cytochrome P450 enzymes involved in deltamethrin metabolism and thus to be an effective synergist with deltamethrin against resistant insects.^{54–58)} PBO has been used at 1 mg/mL, a non-lethal concentration after 48 and 72 hr ingestion (Supplementary Fig. 3). As expected, at 48 hr, the mortality rates were significantly different between deltamethrin/indoxacarb association ($7 \pm 5\%$) and deltamethrin-LNCs/indoxacarb ($34 \pm 5\%$ with $U(4,5)=3.00$, $p=0.0341$) or deltamethrin/indoxacarb/PBO ($32 \pm 7\%$ with $U(4,6)=2.00$, $p=0.0124$) but no significant difference was obtained between mortality rates of deltamethrin-LNCs/indoxacarb and deltamethrin/indoxacarb/PBO (Fig. 7). At 72 hr, the mortality rates were significantly different between deltamethrin/indoxacarb and deltamethrin-LNCs/indoxacarb ($94 \pm 3\%$ with $U(4,5)=0.00$, $p=0.0050$) or deltamethrin/indoxacarb/PBO ($55 \pm 3\%$ with $U(4,6)=0.00$, $p=0.0080$). These increased mortality rates are due to the protection of deltamethrin from detoxification enzymes by LNCs or PBO. However, a significant difference was observed between mortality rates of deltamethrin-LNCs/indoxacarb and deltamethrin/indoxacarb/PBO ($U(5,6)=0.00$, $p=0.0043$). Our data indicated that encapsulation of deltamethrin allows a better protection from detoxification enzymes and could be an effective alternative to PBO to optimize insecticides efficacy and reduce their doses. Furthermore, the deltamethrin-LNCs/indoxacarb synergistic association represents a promising strategy against resistant pest populations which overproduce detoxification enzymes.

Conclusion

In summary, this study demonstrated that deltamethrin-LNCs could be used with indoxacarb to obtain an efficient synergistic association against *A. pisum* larvae. Nanoencapsulation represents an alternative to PBO and an interesting strategy to reduce doses of insecticide in pest management. This formulation has shown high stability *in vitro* against variation of pH, tempera-




Fig. 7. Comparison of indoxacarb toxicity associated with deltamethrin-LNCs or PBO/deltamethrin on *A. pisum* larvae. Comparative mortality rates produced by deltamethrin 10^{-7} M/indoxacarb 3×10^{-6} M, deltamethrin 10^{-7} M/indoxacarb 3×10^{-6} M associated with PBO at 1 mg/mL and deltamethrin-LNCs 10^{-7} M/indoxacarb 3×10^{-6} M at 48 hr and 72 hr. Data are means \pm S.E.M. ($n=4-6$). ns, not significant $p>0.05$; * $p<0.05$; ** $p<0.01$.

ture or light, reflecting potential for a successful applicability in field. Nevertheless, complementary approved analyses should be performed in the future to use such nanopesticides in agriculture. Indeed the high stability of this type of formulation could induce negative effects regarding non-target organisms and a long persistence in environment. This issue will have to be checked in the final development of environmentally friendly nanopesticides.

Acknowledgements

This work was supported by University of Angers for the project ACR 2019 SynPuce. Authors thank Aurélie Rolland and Fabienne Simoneau from the “Plateau Technique Mutualisé Imagerie Cellulaire” (IMAC, SFR QUASAV, Angers, France) for supporting microscopy experiments. Authors also thank Véronique Jean, Julie-Anne Hugel, Eléonore Moreau, Jérôme Bejaud, Marie-Anne Jourdain, Emilie Boulter, Salomé Rousier for their technical assistance.

Declaration of Competing Interest

The authors declare that there is no competing interest.

Electronic supplementary materials

The online version of this article contains supplementary materials, which are available at <https://www.jstage.jst.go.jp/browse/jpestics/>.

References

- <https://www.fao.org/news/story/en/item/1402920/icode/> (Accessed 5 Sept., 2022).
- C. J. A. Bradshaw, B. Leroy, C. Bellard, D. Roiz, C. Albert, A. Fournier, M. Barbet-Massin, J.-M. Salles, F. Simard and F. Courchamp: Massive yet grossly underestimated global costs of invasive insects. *Nat. Commun.* **7**, 12986 (2016).
- R. K. Sandhi and G. V. P. Reddy: Biology, ecology, and management strategies for pea aphid (Hemiptera: Aphididae) in pulse crops. *J. Integr. Pest Manag.* **11**, 18 (2020).
- C. A. Dedryver, A. Le Ralec and F. Fabre: The conflicting relationships between aphids and men: A review of aphid damage and control strategies. *C. R. Biol.* **333**, 539–553 (2010).
- W. H. Jayasinghe, M. S. Akhtar, K. Nakahara and M. N. Maruthi: Effect of aphid biology and morphology on plant virus transmission. *Pest Manag. Sci.* **78**, 416–427 (2022).
- S. Yan, W. X. Wang and J. Shen: Reproductive polyphenism and its advantages in aphids: Switching between sexual and asexual reproduction. *J. Integr. Agric.* **19**, 1447–1457 (2020).
- J.-C. Simon and J. Peccoud: Rapid evolution of aphid pests in agricultural environments. *Curr. Opin. Insect Sci.* **26**, 17–24 (2018).
- J.-C. Simon, S. Stoeckel and D. Tagu: Evolutionary and functional insights into reproductive strategies of aphids. *C. R. Biol.* **333**, 488–496 (2010).
- K. Luo, H. Zhao, X. Wang and Z. Kang: Prevalent pest management strategies for grain aphids: Opportunities and challenges. *Front. Plant Sci.* **12**, 790919 (2022).
- <https://www.fao.org/agriculture/crops/thematic-sitemap/theme/pests/ipm/en/> (Accessed 20 Sept., 2022).
- http://www.endure-network.eu/what_is_endure (Accessed 20 Sept., 2022).
- <https://www.pan-europe.info/> (Accessed 20 Sept., 2022).
- J. P. Deguine, J. N. Aubertot, R. J. Flor, F. Lescourret, K. A. G. Wyckhuys and A. Ratnadass: Integrated pest management: Good intentions, hard realities. A review. *Agron. Sustain. Dev.* **41**, 38 (2021).
- M. Tudi, H. Li, H. Li, L. Wang, J. Lyu, L. Yang, S. Tong, Q. J. Yu, H. D. Ruan, A. Atabila, D. T. Phung, R. Sadler and D. Connell: Exposure routes and health risks associated with pesticide application. *Toxics* **10**, 335 (2022).
- S. Mostafalou and M. Abdollahi: Pesticides: An update of human exposure and toxicity. *Arch. Toxicol.* **91**, 549–599 (2017).
- F. Sanchez-Bayo: Indirect effect of pesticides on insects and other arthropods. *Toxics* **9**, 177 (2021).
- J. Pitti Caballero, L. Murillo, O. List, G. Bastiat, A. Flochlay-Sigognault, F. Guerino, C. Lefrançois, N. Lautram, B. Lapied and V. Aparaire-Marchais: Nanoencapsulated deltamethrin as synergistic agent potentiates insecticide effect of indoxacarb through an unusual neuronal calcium-dependent mechanism. *Pestic. Biochem. Physiol.* **157**, 1–12 (2019).
- K. D. Wing, M. Sacher, Y. Kagaya, Y. Tsurubuchi, L. Mulderig, M. Connair and M. Schnee: Bioactivation and mode of action of the oxadiazine indoxacarb in insects. *Crop Prot.* **19**, 537–545 (2000).
- K. D. Wing, M. E. Schnee, M. Sacher and M. Connair: A novel oxadiazine insecticide is bioactivated in lepidopteran larvae. *Arch. Insect Biochem. Physiol.* **37**, 91–103 (1998).
- https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=&ved=2ahUKewiGk6b3k7L6AhXaQfEDHd1xBPAQFnoECBIQAQ&url=https%3A%2F%2Firac-online.org%2Fdocuments%2Fmoa-classification%2F&usq=AOvVaw2zGC1AY_Ce4U3FEarkHq3t (Accessed 26 Sept., 2022).
- B. Lapied, F. Grolleau and D. B. Sattelle: Indoxacarb, an oxadiazine insecticide, blocks insect neuronal sodium channels. *Br. J. Pharmacol.* **132**, 587–595 (2001).
- K. Silver, K. Dong and B. S. Zhorov: Molecular mechanism of action and selectivity of sodium channel blocker insecticides. *Curr. Med. Chem.* **24**, 2912–2924 (2017).
- M. G. da Silva Oliveira, R. H. Serikawa, F. M. A. da Silva and O. A. Fernandes: Susceptibility and residual effect of indoxacarb formulations on major soybean lepidopteran pests. *Phytoparasitica* **49**, 407–416 (2021).
- A. Maiza, N. Aribi, G. Smagghe, S. Kilani-Morakchi, M. Bendjedid and N. Soltani: Sublethal effects on reproduction and biomarkers by spinosad and indoxacarb in cockroaches *Blattella germanica*. *Bull. Insectol.* **66**, 11–20 (2013).
- J. P. Michaud and A. K. Grant: IPM-compatibility of foliar insecticides for citrus: Indices derived from toxicity to beneficial insects from four orders. *J. Insect Sci. (Ludhiana)* **3**, 18 (2003).
- M. A. Gesraha and A. R. Ebeid: Impact of indoxacarb and sulphur formulation on aphid and three specific predators in Okra fields. *Bull. Natl. Res. Cent.* **45**, 10 (2021).
- Y. Zuo, K. Wang, F. Lin, Y. Li, X. Peng, J. C. Piñero and M. Chen: Sublethal effects of indoxacarb and beta-cypermethrin on *Rhopalosiphum padi* (Hemiptera: Aphididae) under laboratory conditions. *Fla. Entomol.* **99**, 445–450 (2016).
- International Aphid Genomics: Genome sequence of the pea aphid *Acyrtosiphon pisum*. *PLoS Biol.* **8**, e1000313 (2010).
- A. Sadeghi, E. J. Van Damme and G. Smagghe: Evaluation of the susceptibility of the pea aphid, *Acyrtosiphon pisum*, to a selection of novel biorational insecticides using an artificial diet. *J. Insect Sci.* **9**, 1–8 (2009).
- E. Taillebois, A. Beloula, S. Quinchard, S. Jaubert-Possamai, A. Daquin, D. Servent, D. Tagu, S. H. Thany and H. Tricoire-Leignel: Neonicotinoid binding, toxicity and expression of nicotinic acetylcholine

- receptor subunits in the aphid *Acyrtosiphon pisum*. *PLoS One* **9**, e96669 (2014).
- 31) C. F. Henderson and E. W. Tilton: Tests with acaricides against the brown wheat Mite12. *J. Econ. Entomol.* **48**, 157–161 (1955).
 - 32) J. B. Belden, R. J. Gilliom and M. J. Lydy: How well can we predict the toxicity of pesticide mixtures to aquatic life? *Integr. Environ. Assess. Manag.* **3**, 364–372 (2007).
 - 33) B. Heurtault, P. Saulnier, B. Pech, J. E. Proust and J. P. Benoit: A novel phase inversion-based process for the preparation of lipid nanocarriers. *Pharm. Res.* **19**, 875–880 (2002).
 - 34) G. Bastiat, C. O. Pritz, C. Roider, F. Fouchet, E. Lignières, A. Jesacher, R. Glueckert, M. Ritsch-Marte, A. Schrott-Fischer, P. Saulnier and J. P. Benoit: A new tool to ensure the fluorescent dye labeling stability of nanocarriers: A real challenge for fluorescence imaging. *J. Control. Release* **170**, 334–342 (2013).
 - 35) E. Roger, J. C. Gimel, C. Bensley, A. S. Klymchenko and J. P. Benoit: Lipid nanocapsules maintain full integrity after crossing a human intestinal epithelium model. *J. Control. Release* **253**, 11–18 (2017).
 - 36) T. Chen, B. He, J. Tao, Y. He, H. Deng, X. Wang and Y. Zheng: Application of Förster Resonance Energy Transfer (FRET) technique to elucidate intracellular and *in vivo* biofate of nanomedicines. *Adv. Drug Deliv. Rev.* **143**, 177–205 (2019).
 - 37) E. Taillebois, Z. Alamiddine, C. Brazier, J. Graton, A. D. Laurent, S. H. Thany and J. Y. Le Questel: Molecular features and toxicological properties of four common pesticides, acetamiprid, deltamethrin, chlorpyrifos and fipronil. *Bioorg. Med. Chem.* **23**, 1540–1550 (2015).
 - 38) M. Ye, B. Nayak, L. Xiong, C. Xie, Y. Dong, M. You, Z. Yuchi and S. You: The role of insect cytochrome P450s in mediating insecticide resistance. *Agriculture* **12**, 53 (2022).
 - 39) M. E. A. Elzaki, M. A. Miah, Y. Peng, H. Zhang, L. Jiang, M. Wu and Z. Han: Deltamethrin is metabolized by CYP6FU1, a cytochrome P450 associated with pyrethroid resistance, in *Laodelphax striatellus*. *Pest Manag. Sci.* **74**, 1265–1271 (2018).
 - 40) V. D. Rajput, A. Singh, T. Minkina, S. Rawat, S. Mandzhieva, S. Sushkova, V. Shuvaeva, O. Nazarenko, P. Rajput, Komariah, K. K. Verma, A. K. Singh, M. Rao and S. K. Upadhyay: Nano-enabled products: Challenges and opportunities for sustainable agriculture. *Plants* **10**, 2727 (2021).
 - 41) B. Huang, F. Chen, Y. Shen, K. Qian, Y. Wang, C. Sun, X. Zhao, B. Cui, F. Gao, Z. Zeng and H. Cui: Advances in targeted pesticides with environmentally responsive controlled release by nanotechnology. *Nanomaterials (Basel)* **8**, 102 (2018).
 - 42) D. Wang, N. B. Saleh, A. Byro, R. Zepp, E. Sahle-Demessie, T. P. Luxton, K. T. Ho, R. M. Burgess, M. Flury, J. C. White and C. Su: Nano-enabled pesticides for sustainable agriculture and global food security. *Nat. Nanotechnol.* **17**, 347–360 (2022).
 - 43) M. Chaud, E. B. Souto, A. Zielinska, P. Severino, F. Batain, J. Oliveira-Junior and T. Alves: Nanopesticides in agriculture: Benefits and challenge in agricultural productivity, toxicological risks to human health and environment. *Toxics* **9**, 131 (2021).
 - 44) I. Mubeen, M. F. B. Fawzi Bani Mfarrej, Z. Razaq, S. Iqbal, S. A. H. Naqvi, F. Hakim, W. F. A. Mosa, M. Moustafa, Y. Fang and B. Li: Nanopesticides in comparison with agrochemicals: Outlook and future prospects for sustainable agriculture. *Plant Physiol. Biochem.* **198**, 107670 (2023).
 - 45) X. Zhao, H. Cui, Y. Wang, C. Sun, B. Cui and Z. Zeng: Development strategies and prospects of nano-based smart pesticide formulation. *J. Agric. Food Chem.* **66**, 6504–6512 (2018).
 - 46) Y. Yang, J. Cheng, V. M. Garamus, N. Li and A. Zou: Preparation of an environmentally friendly formulation of the insecticide nicotine hydrochloride through encapsulation in chitosan/tripolyphosphate nanoparticles. *J. Agric. Food Chem.* **66**, 1067–1074 (2018).
 - 47) A. J. Shakesby, I. S. Wallace, H. V. Isaacs, J. Pritchard, D. M. Roberts and A. E. Douglas: A water-specific aquaporin involved in aphid osmoregulation. *Insect Biochem. Mol. Biol.* **39**, 1–10 (2009).
 - 48) E. Mróz, D. Kertowska, A. Nowińska, B. Baran, P. Węgierek and Ł. Depa: Morphological description of the alimentary tract of *Geocautricularia* (Passerini, 1856) (Insecta, Hemiptera, Eriosomatinae). *Zoomorphology* **135**, 305–312 (2016).
 - 49) E. Taillebois and S. H. Thany: The differential effect of low-dose mixtures of four pesticides on the pea aphid *Acyrtosiphon pisum*. *Insects* **7**, 53 (2016).
 - 50) Y. L. Phyu, C. G. Palmer, M. St. J. Warne, G. C. Hose, J. C. Chapman and R. P. Lim: A comparison of mixture toxicity assessment: Examining the chronic toxicity of atrazine, permethrin and chlorothalonil in mixtures to *Ceriodaphnia cf. dubia*. *Chemosphere* **85**, 1568–1573 (2011).
 - 51) A. Abd-Ella, M. Stankiewicz, K. Mikulska, W. Nowak, C. Penner, M. Goulu, C. Fruchart-Gaillard, P. Licznar, V. Afaire-Marchais, O. List, V. Corbel, D. Servent and B. Lapied: The repellent DEET potentiates carbamate effects *via* insect muscarinic receptor interactions: An alternative strategy to control insect vector-borne diseases. *PLoS One* **10**, e0126406 (2015).
 - 52) M. Jankowska, J. Wiśniewska, Ł. Fałtynowicz, B. Lapied and M. Stankiewicz: Menthol increases bendiocarb efficacy through activation of octopamine receptors and protein kinase A. *Molecules* **24**, 3775 (2019).
 - 53) E. Moreau, K. Mikulska-Ruminska, M. Goulu, S. Perrier, C. Deshayes, M. Stankiewicz, V. Afaire-Marchais, W. Nowak and B. Lapied: Orthosteric muscarinic receptor activation by the insect repellent IR3535 opens new prospects in insecticide-based vector control. *Sci. Rep.* **10**, 6842 (2020).
 - 54) M. Ahmad, M. A. Saleem and A. H. Sayyed: Efficacy of insecticide mixtures against pyrethroid- and organophosphate-resistant populations of *Spodoptera litura* (Lepidoptera: Noctuidae). *Pest Manag. Sci.* **65**, 266–274 (2009).
 - 55) D. A. Awan, F. Ahmad, M. A. Saleem and A. R. Shakoori: Synergistic effect of piperonyl butoxide and emamectin benzoate on enzymatic activities in resistant populations of red flour beetle, *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *Environ. Sci. Pollut. Res. Int.* **26**, 14200–14213 (2019).
 - 56) M. R. Fakoorziba, F. Eghbal and V. A. Vijayan: Synergist efficacy of piperonyl butoxide with deltamethrin as pyrethroid insecticide on *Culex tritaeniorhynchus* (Diptera: Culicidae) and other mosquito species. *Environ. Toxicol.* **24**, 19–24 (2009).
 - 57) M. Kongmee, K. Thanispong, S. Sathantriphop, C. Sukkanon, M. J. Bangs and T. Chareonviriyaphap: Enhanced mortality in deltamethrin-resistant *Aedes aegypti* in Thailand using a piperonyl butoxide synergist. *Acta Trop.* **189**, 76–83 (2019).
 - 58) E. Tossou, G. Tapa-Yotto, O. K. D. Kpindou, R. Sandeu, B. Datinon, F. Zeukeng, R. Akoton, G. M. Tchigossou, I. Djègbè, J. Vontas, T. Martin, C. Wondji, M. Tamò, A. H. Bokanon-Ganta and R. Djouaka: Susceptibility profiles of *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) to deltamethrin reveal a contrast between the northern and the southern benin. *Int. J. Environ. Res. Public Health* **16**, 1882 (2019).